

**Directions for use:**

1 tablet once or twice daily to help promote intestinal health†.  
Dissolve in mouth or chew.

CAUTION: Diff-Stat® should be taken 1 hour before or 2 hours after antifungal medication.

Do not use if you have a yeast allergy

Keep bottle tightly closed. Store in a cool, dry place.

Do not expose to excessive heat. No refrigeration required.  
Room temperature stable.

Manufactured in an NSF® GMP registered facility.

**Patent Applied For.**

† These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN.


Item # 80001

## Prebiotic & Probiotic Dietary Supplement

# Diff-Stat®

With Syntrinor™

Helps promote intestinal health†



**Pineapple/Orange**  
**Natural & Artificial Flavors**  
30 CHEWABLE TABLETS

### SUPPLEMENT FACTS

Serving Size: 1 chewable tablet

Servings per Container: 30

Amount Per Serving	% Daily Value	
Calories	10	
Total Carbohydrate	3 g	1%*
Sugars	3 g	**
Syntrinor™	942 mg	**
FOS, (Fructo-oligosaccharides), Saccharomyces boulardii, Bacillus coagulans (includes 8.5 billion CFU)		

\*Percent Daily Values are based on a 2,000-calorie diet.

\*\*Daily Value not established.

**Other ingredients:** Sugar, Dextrose, Hydrogenated Cottonseed Oil, Natural & Artificial Orange flavor, Microcrystalline Cellulose, Natural & Artificial Pineapple flavor, Magnesium Stearate, Silicon Dioxide, Ethyl Maltol.

Formulated and marketed by:

**Medical Nutrition USA, Inc.**

Englewood, NJ 07631 USA

1-800-221-0308 [www.diff-stat.info](http://www.diff-stat.info)

Copyright ©2010 Medical Nutrition USA, Inc.

0 26974 41080 5