

Fiber-Stat[®]

Real People, Real Results...

Fiber-Stat[®] Found to Provide Benefits for Bowel Movement Irregularity and Laxative Use.

An observational study of bowel movement irregularity among long-term care residents showed that Fiber-Stat, a liquid fiber supplement, improved bowel movement regularity, thereby reducing the need for laxatives.

STUDY SIGNIFICANCE

Constipation is the most common digestive disorder in the US, resulting in approximately 2.5 million doctor visits and over \$400 million in cost for laxatives. An estimated 75% of elderly institutionalized patients and nursing home residents are given laxatives as a primary treatment for chronic constipation. Data show that frequent use of laxatives is costly, detrimental to resident comfort and can lead to habituation. This study demonstrates that the use of a low-volume, soluble fiber supplement with FOS results in increased bowel regularity and a significant reduction in the use of laxatives in long-term care residents.

(Study Abstract appears on other side.)

THE CHALLENGE OF CONSTIPATION

Constipation is caused by such factors as a lower fiber diet, dehydration or reduced liquid intake, lack of exercise, laxative abuse, changes in routine, medications and specific diseases. Despite the fact that a regular diet provides enough fiber, the condition must often be treated by a variety of laxatives, including stimulants, bulking agents, powdered fibers and mechanical enemas.

Laxatives can be habit-forming. The colon can begin to rely on laxatives to bring on bowel movements. Over time, laxatives can damage nerve cells in the colon and interfere with the colon's natural ability to contract. People who habitually take laxatives become dependent upon them and may require increasing dosages until, finally, the intestine becomes insensitive and fails to work properly.

Chronic use of laxatives in elderly persons has been associated with numerous complications, including diarrhea, fecal soiling, hypoalbuminemia, and high serum levels of magnesium and phosphorus.¹ In one study of long-term care residents, 50% of nursing home residents were found to be regular laxative users, taking more than 30 doses in a one-month period; 17% took between one and 30 doses per month; and 33% took no laxatives.²

References

1. Nutrition Research Newsletter
2. Human Resource and Dietetics

Medical Nutrition USA[™]

Positive Clinical Outcomes Through Evidence-Based Research

Call 1-800-221-0308 for sampling and information on Pro-Stat[®]

Medical Nutrition USA, Inc. Englewood, New Jersey www.pro-stat.info Copyright ©2006 Medical Nutrition USA, Inc.



A Study Assessing the Benefits of Fiber-Stat® on Bowel Movement Irregularity and Laxative Use

STUDY ABSTRACT

OBJECTIVE:

Assess the therapeutic benefits of Fiber-Stat treatment on bowel movement irregularity and the use of laxatives in long-term residents

MATERIALS AND METHODS:

A total of 37 residents of Bergen Regional Medical Center in Paramus, New Jersey's largest hospital/nursing home complex were enrolled in the study. All residents received 30 ml of Fiber-Stat (P.O. or via GT) twice daily for 60 days. Records on resident's daily bowel movements and their use of laxatives were maintained and analyzed.

RESULTS:

There were a total of 89 standing laxatives ordered by the 37 residents during the observational period. At the onset of the study, prior to taking Fiber-Stat, there were a total of:

- 65 standing laxative orders (74%)
- 24 PRN (as needed) laxative orders (26%)

Changes in laxative orders following Fiber-Stat therapy included:

- 49 standing laxative orders
- 19 PRN laxative orders

There was an overall 25% reduction in laxative orders (from 65 to 49) following Fiber-Stat


There was an overall 21% reduction in PRN laxative orders (from 24 to 19) following Fiber-Stat

A cumulative 24% reduction in laxative and PRN orders over a 60 day period following Fiber-Stat supplementation.

CONCLUSIONS

- Fiber-Stat reduced the dependence on laxatives among this patient population.
- Fiber-Stat helps regularity in bowel movements of long-term care patients.

Figure. Bergen Regional Medical Center, Paramus, NJ, Long-Term Care Complex Change in Laxative Use Following a 60-day Fiber-Stat Therapy



Key Finding:
Fiber-Stat® had a cumulative 24% reduction in laxative usage & PRN orders over a 60 day period following Fiber-Stat supplementation among elderly long-term care residents.

Sponsored by:



MEDICAL NUTRITION USA, Inc.

For a copy of the complete study, call 201-569-1188 or write to Medical Nutrition USA Inc., 10 West Forest Avenue, Englewood, NJ 07631